



SURF LIFE SAVING®
NEW ZEALAND

Health, Safety and Welfare Bulletin

Date: 30 August 2019

To: All SLSNZ Clubs

From: Ross Merrett – Health, Safety, Welfare and Risk Manager

Subject: SLSNZ's Response to Measles Outbreak

Background

There are increasing incidents of measles throughout New Zealand, particularly in Auckland. Measles is highly contagious. Cases of measles are expected to rise in the foreseeable future, particularly in Auckland. The SLSNZ 2019 Pool Rescue Championships are scheduled for 11-13 October in Auckland, as well as other SLS events around the country as we start to build into the 2019-20 season.

What is the problem?

Those not immunised against measles are at risk of contracting and spreading measles to others, particularly when gathering indoors, and/or for sports events, where athletes are in close contact with others, such as pool rescue events.

What is being done to address the problem?

SLSNZ requests that members who are not yet immunised do not attend the forthcoming 2019 Rescue Pool Championships unless they have received an MMR vaccination at least 2 weeks prior to the event i.e. by 26 Sept. 2019. Likewise those not yet immunised are requested not to attend any other SLS events unless they have completed their MMR vaccinations at least 2 weeks prior to the scheduled event, until further notice.

What next?

Any person who is suspected of having measles must be sent home until they are seen by a doctor. If measles is confirmed, they must remain at home away from any visitors, unless these visitors are known to be immune.

Those who have not been vaccinated, and who have been in a confined space with a confirmed case of measles should stay at home if they are not immune, and check for symptoms until the end of the quarantine period. This might be one to two weeks depending on when they were exposed to measles.

What are the symptoms of measles?

Measles symptoms include a high fever, runny nose, cough and sore red eyes, followed several days later by a rash starting on the face and neck before spreading to the rest of the body. Someone with measles can spread it to others from five days before the rash appears until five days after.

If somebody is confirmed as having measles, they will need to stay at home in isolation until five days after their rash first appeared.

It's important to check you are up to date with your immunisations, especially if you're travelling overseas. For more information please seek advice from the [Ministry of Health web site](#).

In it for life



FINANCIAL SERVICES



Surf Life Saving New Zealand

2 Boeing Place, Mount Maunganui 3116. PO Box 4667, Mt Maunganui 3149. www.surflifesaving.org.nz